

SKULARI, PETR

CZECH

Application of microstructure analysis for technology of
plastic magnesium alloys. Petr Skulari (Výzkumný ústav
kovů, Pauderské Úřezny, ~~Czechoslovakia~~ ~~Mnichův Lázeň 10,~~
200-15(1955).—The advantages of microstructure analysis
in the technology of plastic Mg alloys is pointed out and
the influence of hexagonal lattice on the mech. and tech-
nological characteristics of these alloys is shown.
Petr Schneider

M

A RSK

SKULARI PETR

3

CZECH 11917* Micro-Examination of Dynamic Fatigue. Sledování dynamické únavy mikrorozsvětlením. (Czech) Petr Skuláří and Vladimír Obdržálek. Hutnické Listy, v. 10, no. 3, May 1955, p. 279-284.
Suppositions for the practical use of the examination and results obtained for Al and its alloys under alternating dynamic loads. Explains the physical principles of fatigue related to material in annealed and hardened conditions. Graphs, micrographs, diagrams, table. 9 ref.

[Handwritten signature]

SKULARI, F.

Some properties of semifinished duralumin products and their heat treatment.

p. 949 (Hutnicke Listy) Vol. 12, no. 10, Oct. 1957, Praha, Czechoslovakia

SO: MONTHLY INDEX OF EAST EUROPEAN ACCESSIONS (EEAI) LC, VOL. 7, NO. 1, JAN. 1958

*Distr: AE2c 27

V Röntgenography of titanium. Petr Skulari and Ludmila
Charvatova. *Hutnické listy* 10, 333-334 (1958). Based
upon systematic examn. of Ti structure with microröntgen
methods, the effect was detd. of different elements on type
and dimensions of lattice of α - and β -Ti. They detd. also
the various effect of different elements on the deformation
of Ti lattice and at the same time on its plastical deforma-
tion. O, Ni, H, C, and Si are supposed to give the most in-
terference. O and N cannot be removed from the lattice
even with vacuum annealing. Petr Schneider

5
1

17 JG

67099

18.12.10

AUTHORS: Skulari, Petr, Doctor and Chvátalová, Ludmila
CZECH/34-59-12-2/44
TITLE: X-ray Investigation¹ on the Hardening of Special Silumins
PERIODICAL: Hutnické listy, 1959, Nr 12, pp 1032 - 1038

ABSTRACT: A literary search has revealed the possibility of improving the mechanical properties of certain alloys by correct ageing but has revealed no detailed information on the conditions of ageing (temperatures of hardening and tempering and the appropriate heating times) and on the suitable composition of special silumins. It appears that so far the kinetics of unsaturated solid solutions of Al alloys, which changes with the addition of small quantities of certain elements, has not been adequately studied. Neither has the problem of the suitable composition of such special silumins been studied. The authors of this paper have studied the influence of small quantities of Mn and Ti on the kinetics of ageing of special silumins, containing 5% silicon, as a function of the conditions of the artificial ageing and the contents

4

Card 1/5

67099

CZECH/34-59-12-2/44

X-ray Investigation on the Hardening of Special Silumins

of a given addition. The changes in the structure during the shaping and the decomposition of the saturated AlSi solid solution were studied by X-ray structural analysis which provides the possibility of recording not only the character of the structure (size of the crystals, phase distribution) but also, by limiting the parameters, determining the state of the solid solution, disturbances in the lattice and the internal crystalline structure of the alloy. The structural studies were supplemented by hardness tests. The aim of the authors was to determine a suitable element, the required quantity of this element and the optimum conditions of artificial ageing (which can be achieved by short-duration heating prior to hardening and tempering) so as to produce an alloy with better mechanical properties. The authors assume that the prevailing view that decomposition of alloying elements in the solid solution of Al lasts a relatively long time is based on incomplete knowledge of the influence of small additions on the kinetics of saturated solid solutions. For producing the experimental alloys, the authors used

Card2/5

67099

CZECH/34-59-12-2/44

X-ray investigation on the Hardening of Special Silumins

a 99.5% purity Al, silicon containing 0.57% Fe, cathodic copper, electrolytic manganese, 99.90% purity Mg and Ti produced by the Kroll method. The heats were produced by high-frequency smelting or in an electric resistance furnace with a graphite crucible of about 1 kg capacity. The charge was fed in jointly with Mn and Cu and after smelting the melt was covered with salts of eutectic composition. Further alloying was effected in the following sequence: Si; Ti; Mg. Mg was added in the form of AlMg 10% and Ti in the form of AlTi 5%. The alloy was cast at 750 °C into small unpainted ingot moulds. The authors discuss in some detail the results obtained relating to the influence of Cu, Mg, Mn and Ti on the hardening of the investigated silumin alloys. In the preliminary tests, it was found that addition of 0.2% Mn to the base alloy containing 5% Si, 0.2% Cu and 0.25% Mg brought about a considerable refining of the structure, which was very uniform. In the further tests, the influences of Mn and Ti were investigated in greater detail

Card 3/5

67099

CZECH/34-59-12-2/44

X-ray Investigation on the Hardening of Special Silumins

on two alloys which contained 5% Si and 0.2% Cu and differed in their Mg and Mn contents which were, respectively, 0.42 and 0.53% Mg and 0.33 and 0.24% Mn. It was found that addition of 0.2% Mn or 0.2% Ti leads to a considerable refining of the structure of the alloy and to an increase in the hardness of special silumins, containing 5% Si, after hardening. For the alloy containing 5% Si, 0.2% Cu, 0.5% Mg, 0.14% Fe, 0.2% Mn, rest Al, the optimum heat-treatment conditions are: quenching in cold water from 450 °C after a heating time of 30 - 120 min; tempering from 180 °C after a heating time of 120 - 240 min. For an alloy of equal composition but containing 0.2-0.3% Ti instead of 0.2% Mn, the optimum heat-treatment conditions are: quenching in cold water from 500 °C after heating for 120 - 140 min; tempering from 180 °C after heating for 120 - 140 min. These heat-treatment conditions ensure a uniform structure of fine crystals, a constant lattice volume and an increase in the hardness by 100% as compared with that in the as-cast state; thereby Ti proved more active than Mn. The X-ray

Card4/5

£7099

CZECH/34-59-12-2/44

X-ray Investigation on the Hardening of Special Silumins

method used permits determining the trend for the further development of $AlSi_5$ alloys and proposing a suitable

composition; the tests require a very small quantity of material and very simple preparation of the specimens.

However, these tests have to be supplemented by foundry experiments and mechanical tests.

There are 7 figures and 13 references, of which 4 are Czech, 3 German, 2 English and 4 Soviet.

ASSOCIATION: Vyzkumný ústav kovů, Panenské Břežany
(Metal Research Institute, Panenské Břežany)

SUBMITTED: February 7, 1959

4

Card 5/5

S/137/62/000/011/033/045
A005/A101

AUTHOR: Skylari, Peter

TITLE: X-ray-diffraction phase analysis of metals at high temperatures

PERIODICAL: Referativnyy zhurnal, Metallurgiya, no. 11, 1962, 94, abstract
111628 ("Kohász. lapok", 1962, v. 95, no. 1, 40 - 45, Hungarian)

TEXT: A long duration of the process and the lack of a chamber to analyze the whole production process, are the deficiencies of the X-ray diffraction phase analysis. The existing chambers show only the changes in the intensity of the lines, but the metal composition can not be determined. Information is given on measurements in a special chamber of the integrated intensity of two diffraction lines with the use of two Geiger-Muller counters. In quantitative phase analyses of the steel and Ti the specimen under investigation of 12 x 20 x 3 mm in size was heated within some seconds to 1,000 - 1,500°C by direct passage of 100 amps current. The temperature was measured with two Pt/Pt-Rh thermocouples. A vacuum of $1 \cdot 10^{-5}$ - $1 \cdot 10^{-6}$ mm Hg was produced in the chamber. With the use of this chamber the following studies can be performed: 1) quantitative phase ana-

Card 1/2

X-ray-diffraction phase analysis of...

S/137/62/000/011/033/045
A006/A101

lysis of steel and alloys in a wide temperature range; 2) analysis of diagrams of stable and metastable states; 3) investigation of metal transition points from one phase condition to another; 4) investigation of quenching and tempering processes during heating and cooling. The chamber can also be used to investigate Ti and its alloys.

B. Mat'ush

[Abstracter's note: Complete translation]

Card 2/2

KOREJTKO, J.; SELVEK, A.; SKULAVIK, L.

Porosity of the grain mass. Zemedel' tech 11 no.2:101-106 F '65.

1. Chair of Physics of the Faculty of Operation and Economic
of the Higher School of Agriculture, Nitra. Submitted July 27,
1964.

VISHNEVSKIY, Palladiy Fedorovich [Vyshnevs'kyi, P.F.];
SKUL'BASHEVSKIY, A.D. [Skul'bashevs'kyi, A.D.], inzh.,
retsensent; BILASH, I.K., red.

[Showers and shower runoff in the Ukraine] Zlyvy i zly-
vovyi stik na Ukraini. Kyiv, Naukova dumka, 1964. 288 p.
(MIRA 17:8)

KIRKEVICH, G.A.; SKUIENKO, A.I.

Making use of experience abroad in the construction of gas
reservoirs in the Ukraine. Neft. i gaz. prom. no.2:65-68
Ap-Je '63. (MIRA 17:11)

SKULIC, M.

SKULIC, M. Protective equipment in the high-tension network of Bosnia
and Hercegovina. p. 216

Vol. 9, no. 4/5, Apr./May 1956

ELEKTROPRIVERDA

TECHNOLOGY

Beograd

So: East European Accession, Vol. 6, no.3, March, 1957

SKULIC, Mirko, inz.

The Yugoslav-made 170/280-K³ type suspension insulators
in the transmission network of Yugoslavia. Elektroprivreda
15 no.6/7:315-320 Je-J1 '62.

1. Elektroistok, Beograd.

SKULIMOVSKAYA, G.A.

Control of microsporia. Sovet. med. 16 no. 6:39-41 June 1952.
(CLML 22:4)

1. Of the Mycological Department (Head -- Prof. A. M. Ariyevich),
Central Skin-Venereological Institute (Director -- Candidate Medical
Sciences N. M. Turanov), Ministry of Public Health USSR.

SKULIMOVSKAYA, G.A. (Moskva)

Local corticosteroid treatment of some skin diseases. Vrach. delo
no.10:122-124 0 '61. (MIRA 14:12)
(CORTICOSTEROIDS) (SKIN—DISEASES)

LEONOVA, V.I., kand.med.nauk; SKULIMOVSKAYA, G.A.

Treatment of chronic eczema of the external ear canal with
corticosteroid hormone ointments. Vest. dermat. i ven. 38
no.4:79-81 Ap '64. (MIRA 18:4)

1. 4-ye Glavnoye upravleniye pri Ministerstva zdravookhraneniya
SSSR (nauchnyy rukovoditel' raboty - chlen-korrespondent SSSR
prof. V.A.Rakhmanov), Moskva.

SKULIMOWSKI, Andrzej (Krakow, Boh. Stalingradu 78 m. 16.)

The oldest royal privileges given to a Polish physician. Polski tygod.
lek. 13 no.47:1895-1896 24 Nov 58.

(HISTORY, MEDICAL
in Poland (Pol))

Janowski, Henryk

Henryk Janowski 1913-1984. Pol. wyg. lek. 19 no.8:312-314
1984.

History of science and teaching of medical history at the
Jagellonian University and Academy of Medicine in Krakow.
Ibid.:315-318

SKULIMOWSKI, Mieczyslaw.

Use of garlic for medical purposes. Polski tygod.lek. 11 no.2:68-69 9 Jan 1956.

1. Z Zakladu Historii Medycyny A.M. w Krakowie. Krakow, ul. Kopernika 7.
Z akład Historii Medycyny A.M.

(VEGETABLES,
garlic, medicinal use)

SKULIMOWSKI, Mieczyslaw

Wiktor Feliks Szokalski about Adam Mickiewicz. Polski
tygod. lek. 12 no.11:409-411 11 Mar 57.

1. (Z Zakladu Historii Medycyny A.M. w Krakowie). Adres:
Wieliczka, ul. A. Grottgera 1.

(BIOGRAPHIES,
Szokalski, Wiktor F. (Pol))

(LITERATURE,
W.F. Szokalski's views on literary works of poet
Mickiewicz (Pol))

SKILL IN 1956, M.

Copernicus, the physician. Orv. hetil. 106 no.30:1423-1426
25 J1'65.

SKULIMOWSKI, Mieczyslaw

Development of concepts of hypertension in Polish medicine.
Polski tygod. lek. 12 no.24:920-924 10 June 57.

1. Z I Kliniki Chorob Wewnętrznych A.M. w Krakowie; kierownik:
prof. dr. Leon Tochowicz i z Zakładu Historii Medycyny A.M. w
Krakowie; kierownik: zast. prof. dr. Zbigniew Kukulski. Adres:
Krakow, ul. Kopernika 17. I Klinika Chorob Wewnętrznych A.M.
(HYPERTENSION, history,
in Poland (Pol))

SKULIMOWSKI, Mieczyslaw

History of the Department of Medicine of Krakow University
prior to the Kollanta; Reform (1364-1779). Pol. tyg.lek. 18
no.50:1899-1902 9 D'63

*

TOCHOWICZ, Leon; SKULIMOWSKI, Mieczyslaw

Development and teaching of internal medicine at the Department
of Medicine of the Jagiello University and Academy of Medicine
in Krakow. Pol. tyg. lek. 18 no.40:1494-1497 30:8:163.

(HISTORY OF MEDICINE) (EDUCATION, MEDICAL)
(INTERNAL MEDICINE) (SCHOOLS, MEDICAL)

POPIELA, Tadeusz; SKULIMOWSKI, Mieczyslaw

History of the School of Medicine of the Jagiello University
during the period of re-Polonization until the liberation
1864-1918). Pol. tyg. lek. 19 no.3:112-113 20 Ja^r64

KANIA, Jerzy, inz.; SKULIMOWSKI, Wojciech, mgr inz.

Control receiver of color TV pictures. Prace Inst teletechn 6
no.2:103-111 '62.

LARYUKHIN, M.A.; REKUNOV, V.S.; SKULINA, K.I.; MIKHINA, L.I.

Use of a fire chemical spray from an airplane in controlling tick-borne encephalitis carriers in the Anzhero-Sudzhensk District in 1957-1959. Med.paraz.i paraz.bol. no.3:347-351 '62. (MIRA 15:9)

1. Iz otdela entomo-toksikologii i dezinseksii (zav. - prof. V.A. Nabokov) Instituta meditsinskoy parazitologii i tropicheskoy meditsiny imeni Ye.N. Martsinovskogo (dir. - prof. P.G. Sergiyev) Ministerstva zdravookhraneniya SSSR, Gosudarstvennogo Nauchno-issledovatel'skogo instituta grazhdanskogo vozdushnogo flota (nach. - general leytenant N.A. Zakharov) i Anzhero-Sudzhenskoy gorodskoy sanitarno-epidemiologicheskoy stantsii.

(ENCEPHALITIS) (AERONAUTICS IN INSECT CONTROL)
(ANZHERO-SUDZHENSK DISTRICT--TICKS AS CARRIERS OF DISEASE)

1. *Journal of the American Medical Association*, 1955, 161, 1000.

1. Study of the East European Accessions, (1941), 11, Vol. 2, no. 10, Oct. 1956, 1961.

SKULJ, A.

SKULJ, A. Constructive development of the high-tension network in Slovenia.
p. 242

Vol. 9, no. 4/5, Apr./May 1956
ELEKTROPRIVERDA
TECHNOLOGY
Beograd

So: East European Accession, Vol. 6, no.3, March, 1957

SKULJ, K.

Yugoslavia (430)

General - Serials

The Slovenian land. p 50. SLOVENSKA BESEDA. (Zalozba Slovenska Beseda)
Buenos Aires. (Monthly illustrated dealing with cultural problems of
Slovenian emigrants in Argentina) Vol. 2, no 3-4, Mar.-Apr. 1951.

East European Accessions List. Library of Congress, Vol 1, no. 13,
November 1952. UNCLASSIFIED.

SKULJ, Vladimir, Dr.

Fourth stage of labor. Lijec. vjes. 78 no.11-12:530-534
Nov-Dec 56.

1. Iz Klinike za zenske bolesti i porode Medicinskog fakulteta
u Zagrebu.

(LABOR

fourth stage (Ser))

SKULJ, Vladimir

Manchester operation. Srpski arh. celok. lek. 85 no.6:688-699 June 57.

1. Klinika za zenske bolesti i porode Medicinskog fakulteta u Zagrebu.

Upravnik: prof. dr. Stjepan Vidakovic.

(UTERUS, dis.

prolapse, surg., Fothergill operation (Ser))

SKULJ, Vladimir, Dr.; STOILJKOVIC, Cedomir, Dr.

On toxoplasmosis in obstetrics. Lijec vjes 82 no.6:501-506 '60.

1. Iz Klinike za zenske bolesti i porode Medicinskog fakulteta
Sveucilista u Zagrebu
(TOXOPLASMOSIS in pregn)
(PREGNANCY compl)

ROSENZWEIG, Erich; BERGER, Milan; SKULJ, Vladimir

Dorso-occipital presentation. Rad. med. fak. Zagreb 9 no.1:37-69 '61.

(LABOR PRESENTATION)

ROSENZWEIG, Erik, dr.; SKULJ, Vladimir, dr.; PAVLIC, Zdravko, dr.

Role of modern general practitioners in obstetrics. Liječn. vjesn.
83 no.11:1171-1175 '61.

1. Iz Klinike za ženske bolesti i porođaje Medicinskog fakulteta
u Zagrebu.

(OBSTETRICS) (GENERAL PRACTICE)

SKULJ, Vladimir

On a possibility of spontaneous cure in tubal pregnancy. Srpski
arh. celok. lek. 89 no.3:275-283 Mr '61.

1. Klinika za zenske bolesti i porode Medicinskog fakulteta Univerzi-
teta u Zagrebu. Upravnik: prof. dr Stjepan Vidakovic.

(PREGNANCY ECTOPIC)

DEKARIS, Mihovil; RAJHVAJN, Branko; SKULJ, Vladimir; DRAZANCIC, Ante

Endometrial carcinomas. Rad. med.fak. Zagreb. 10:201-213 '62.
(UTERINE NEOPLASMS)

S

KARTASHOV, A.K.; KAGANOV, I.N., kand. tekhn. nauk, spets. red.;
SKULKOV, G.S., otv. za vyp.; DMITRIYEVA, Ye.P., otv. za
vyp.; RYBAKOVA, L.G., tekhn. red.

[New methods for the purification of diffuser juices] Novye
metody ochistiki diffuzionnogo soka. Moskva, TSentr. in-t
nauchno-tekhn. informatsii pishchevoi promyshl., 1962. 30 p.
(MIRA 16:4)

(Sugar manufacture)

LOYTSYANSKAYA, M.S.; SKULKOV, G.S., otv.za vyp.; KUDRYAVTSEVA, A.P.,
otv. za vyp.; RYBAKOVA, L.G., tekhn. red.

[Microbiological foundations of the production of vinegar]
Mikrobiologicheskie osnovy proizvodstva uksusa. Moskva,
TSentr. in-t nauchno-tekhn. informatsii pishchevoi pro-
myshl., 1962. 35 p. (MIRA 16:4)
(VINEGAR—MICROBIOLOGY)

JKULKOVA, E.C.

Supervision of the activities of pharmaceutical institutions.
App. delo 13 no.4:69-73 FI-Ag '64. (MIRA 13:3.

1. Tsentral'nyy aptekhnny nauchno-issledovatel'skiy institut,
Moskva.

SAULKOVA, R.O.

Economic councils in pharmaceutical institutions. Apt. 56 6 13
nr. 5:51-53 3-6 1-4. (1981 12:13)

1. Tsentral'nyy aptekhnny nauchno-issledovatel'skiy institut,
Moskva.

PANCHENKO, Ye.I.; SKULKOVA, R.S.

Experience in the organization of interhospital drugstores.
Apt. delo 14 no. 4:8-13 J1-Ag '65 (MIRA 19:1)

1. Tsentral'nyy aptechnyy nauchno-issledovatel'skiy institut,
Moskva. Submitted January 21, 1965.

M. Phys
②
5

Photographic abs

P 4 1953

Industrial + Scientific
Application of Photography

The threshold of electron emission in Ni and Mo is measured for different energies of the incident electrons. The emission curves are given, and the velocity of the emitted electron is compared with the energy of the incident electron. (J. & R.)

THE THRESHOLD OF ELECTRON EMISSION IN NICKEL AND MOLYBDENUM. A. R. Skul'man and E. I. Myshkin. Doklady Akad. Nauk S.S.S.R. 91, 1075-8 (1953) Aug. 11. (In Russian)

6-15-54
RMZ

SKUL'MAN, A. R.

THERMOELECTRON EMISSIONS FROM THIN PLATES OF
THORIUM OXIDE AND THORIUM ON MOLYBDENUM.
A. R. Skul'man and A. P. Rumyshev. Doklady Akad. Nauk
S.S.S.R. 23, 455-8 (1963) Nov. 21. (In Russian)

32

SKULME, K. A.

SKULME, K. A. -- "Investigation of the Casualty and Diet Therapy of Stomach Ulcers -- Duodenum." Latvian State U, 1949. In Latvian
(Dissertation for the Degree of Candidate of Medical Sciences)

SO: Izvestiya Ak. Nauk Latviyskoy, SSR, No. 9, Sept., 1955

1. Skulme, K., Alless, A., Sukute, H.

2. USSR (600)

4. Histamine

7. Latv. PSR Zin. Akad. Vestis no. 12, 1950.

9. Monthly List of Russian Accessions. Library of Congress, March 1953, Unclassified.

ROBEZHNIKES, I.A.; SKULME, K.A.; PETERSON, M.P.

Oxidation of ascorbic acid by human gastric juice. Latvijas PSR Zinātņu
Akad. Vēstis '51, 757-60. (MLRA 5:10)
(CA 47 no.22:12597 '53)

SKULME, K.A., kand.med.nauk, PRANIS, L.Yu.,

Materials on morphologic changes of blood in cancer. Vopr.klin.lech.
zlok. novoobraz., Riga 1:39-50 1953

(NEOPLASMS, diag.

blood tests

(BLOOD, in various dis.

cancer, morphole changes, diag. value

SKULME, K.A., kand.med.nauk

Nutrition in cancer. Vopr.klin.lech.zlok.novoobraz., Riga 1:134-144
1953

(DIETS, in various dis.
cancer
(NEOPLASMS, ther.
diet

SKULME, K.A.

✓The protein composition of blood of cancer patients. K. A. Skulme. *Trudy Inst. Eksp. Med., Akad. Nauk S.S.S.R.* 1953, No. 3, 51-60; *Referat. Zhur. Khim., Biol. Khim.* 1953, No. 7105. —Studies were made of total proteins in 606, albumin and globulin in 430, and fibrinogen in 102 patients with cancer of different body parts. Protein detns. were made refractometrically; fibrinogen by the method of Cullen-Van Slyke; globulin by pptn. with $(NH_4)_2SO_4$. The protein content is related to the site of the cancer, its extent, and the onset of destructive processes. Less than 7% of protein was found in 60% of patients with inoperable forms of cancer of the stomach and 40% of operable patients. Protein content of less than 6% was found resp. in 15 and 10% of cancer patients. In cancer of other organs of digestion the percentages were 2.5 and 17%, resp. In cancer of the skin a protein content of 7% or less occurred in 5.5% of the patients, and in the cancer of the breast no hypoproteinemia was observed. Most pronounced hypoproteinemia was observed in cases with metastases, cachexias, and high erythrocyte sedimentation rates. In patients with ascites, transudation, purulence, prolonged vomiting, etc., the protein concn. remains normal. In patients with gastric and duodenal ulcers serum protein concns. below 6% were found in 1.4% and below 7% in 13%. Except for the cancer of the skin, the albumin-globulin ratio was lowered in 73-88% of cancer patients regardless of site of the neoplasm. The ratio is particularly low in cachexias, chronic illnesses, in metastatic disease, and in the presence of destructive processes. No interrelation was recorded between the erythrocyte sedimentation rate, degree of anemia, and the albumin-globulin ratio. A high fibrinogen content (0.4%) was observed in patients with severe cachexia with destructive processes and with above normal temp. Low values (below 0.2%) were observed in the presence of extensive metastatic lesions and functional liver disturbances. In patients with a satisfactory state of health fibrinogen remained within normal limits (0.2-0.4%). B. S. L.

1. SKULF, K. A.
2. USSR (600)
4. Nutrition - Latvia
7. Conference on problems of nutrition, held by the Departments of Biology and Agriculture of the Academy of Sciences of Latvia in Riga on September 16-17, 1952. Vop. pit. 12, No. 2, 1953.
9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

SKULME, K.A., kand.med.nauk

Comparative studies of gastric function in cancer and precancerous conditions. Vopr.klin.lech.zlok.novoobraz., Riga 2:21-23 1955

1. Sektor onkologii (zav., prof. doktor P.I. Stradyn') Instituta eksperimental'noy meditsiny AN Latvyskoy SSR (dir. prof. doktor P.Ya.Gerke).

(STOMACH, neoplasms
cancer & precancer, comparison of motor funct. (Rus))

SKULME, K.A., kand.med nauk

Nutrition of gastric patients and its relation to surgery. Vopr.
klin.lech.zlok. nevoobraz., Riga. 2:147-173 1955

1. Sektor onkologii (zav. prof. doktor P.I. Stradyn') Instituta
eksperimental'noy meditsiny AN Latvyskoy SSSR (dir. - prof.doktor
P.Ya. Gerke).

(STOMACH, surgery,
diets in (Rus))

(DIETS,
in stomach surg (Rus))

USSR/Medicine - nutrition

FD-3077

Card 1/1 Pub. 141 - 23/23

Author : Skulme, K. A.

Title : Scientific conference on problems in therapeutic nutrition

Periodical : Vop. pit., 61-63, May/Jun 1955

Abstract : A scientific conference on problems in therapeutic nutrition was held in Riga on 18-19 November 1954 under the auspices of the Academy of Sciences Latvian SSR, the Ministry of Health Latvian SSR, and the Riga Medical Institute. Scientists representing the Institute of Nutrition Academy of Sciences USSR and the Military-Naval Medical Academy also attended. Seventeen reports concerning theoretical problems in therapeutic nutrition, vitamins, therapeutic nutrition for a number of illnesses, and nutrition for healthy and sick babies were presented.

Institution :

Submitted :

SKULME, K.

GENERAL

PERIODICALS: VESTIS, NO. 6, 1958

SKULME, K. Significance of research on leukopedesis of the stomach
for the diagnostic of stomach diseases. p. 89.

Monthly List East European Accessions (EEAI) LC, VOL. 8, No. 2,
February 1959, Unclass.

SKULME, Kseniya Andreyevna (Inst of Experimental Med, Acad Sci Latvian SSR) for
Doc Med Sci on the basis of dissertation defended 22 June 59 in Council of Riga
Med Inst, entitled "Problems of the etiology, pathogenesis and early diagnosis
of cancer and precancerous gastric diseases." (BMVSSO USSR, 1-61, 21)

-88-

8/851/62/000/028/014/015
D296/D307

AUTHOR: Skulme, K.A.
TITLE: A rational diet for patients with malignant disease
and receiving radiotherapy
SOURCE: Akademiya nauk Latviyskoy SSR. Institut eksperiment-
tal'noy i klinicheskoy meditsiny. Trudy, no. 28, 1962.
Znachenie faktora pitaniya v profilaktike luchevoj
bolezni, no. 4, 171 - 244

TEXT: 290 patients with malignant disease, mainly women suffering
from carcinoma of the cervix (aged between 41 and 60) received deep
x-ray therapy in a single dose of 250 - 300 r twice daily up to a
total of 18 000 r, or 1 local treatment with Co60 in a single dose
of 1800 - 2600 mg/hr. up to a total of 8000 - 12,000 mg/hr. for 30
- 60 days. The patients were assessed every 5 - 10 days with regard
to their general condition, hemopoietic function, protein and vita-
min metabolism, and the secretory and motor functions of the sto-
mach. The patients were divided into 4 groups, the first of which
(66) was given the usual hospital diet. The second group of 67 recei-
Card 1/3

A rational diet for patients with ...

S/851/62/000/028/014/015
D296/D307

ved a diet enriched with proteins and vitamin C, the third (of 54) was given in addition vitamins B¹, B₂, B₆ and PP. Finally, the 4th group of 32 patients received all these and also pantothenic acid, folic acid and rutin. An earlier investigation showed that the usual hospital diet with an average protein ration of 1 g/kg daily was insufficient to maintain the nitrogen balance of patients with malignant disease and receiving radiotherapy. Experiments were carried out on rats and guinea pigs. General digestive and nervous disorders were observed most frequently, in the 1st and least frequently in the 4th group. Renal disorders, hemorrhages and a reduction in blood pressure occurred in all 4 groups. The immediate results of the treatment were most satisfactory in the 4th group. The state of the protein reserve was investigated in 172 patients by estimation of the whole blood globin level and the total serum proteins. Radiotherapy caused a marked decrease in the blood globin and serum protein levels, most often in the 1st and least of all in the 4th group. A fall in the globin level could also be produced experimentally in rats exposed to total body radiation. Radiation also decreased the vitamin C level in the adrenal cortex and blood.

Card 2/3

A rational diet for patients with ...

S/851/62/000/028/014/015
D296/D307

This fall could only be observed in patients receiving the ordinary hospital diet but in the other 3 groups it was compensated by the high doses of vitamin C which these groups received. As a rule, radiotherapy led to a decrease in the gastric acid secretion but in some hyperacid patients increased gastric secretion could be observed. These disorders of secretory function were most marked in the 1st group and least marked in the 4th group. Changes in the ammonia and lactic acid contents of the gastric juices were less pronounced. The motor function of the stomach was impaired most in the 1st and least in the 4th group. The same applied to the loss of appetite and dyspepsia. The author concludes that patients with malignant disease require in the period of radiotherapy a diet reinforced by proteins and vitamins. There are 27 figures and 38 tables. ✓

Card 3/3

SKULME, K.A.

Scientific session in Leningrad dedicated to the problem of
chronic gastritis. Vop. pit. 22 no.6:84-86 N-D '63.
(MIRA 17:7)

Effect of em and fat on the growth of gastric cancer in
rat. Vop. pit. 21 no. 1: 1988. (MIRA 1840)

Laboratory of pathology, Prof. A. A. Shulze, Latvian
Institute of Experimental Medicine and Biophysics, AN SSSR,
Riga.

CHŁOPCZYŃSKA-ERYKOWSKA, Danuta; LARSKI, Stanisław


Changes in the transaminase activity (GOT and GPT) in the allantoic fluid of chick embryos infected with Newcastle disease virus. Med. dosw. mikrobiol. 16 no.4:87-92 1974.

1. 2 II Prace Instytutu Wirusologii Ogrodnej i W. Polowy (Kierownik: doc. dr. Z. Larski).

SKULNIEWSKA-KRYSZEKOWSKA, Danuta; LARSKI, Zdzislaw

Effect of some antihistaminic drugs on the multiplication of
Newcastle disease virus. Med. dosw. mikrobiol. 17 no.3:251-256
1965.

1. Z II Pracowni Wirusologii Ogolnej Instytutu Weterynarii w
Pulawach (Kierownik: doc. dr. Z. Larski).



1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
PROCESSING AND PROPERTY INDEX																																																			
<p><i>Handwritten: B3</i></p> <p><i>Handwritten: 1</i></p> <p>Dynamometric measurement of work performed by horse. J. Shulmovski (Ann. Univ. M. Curie-Skłodowska, 1960, 8, 2, 1-12).—Work performed in one day by farm horses is classified as light <1.5, medium 1.5-2.5, heavy 2.5-3.5, and very heavy >3.5 million kg.-m. R. Tauson.</p>																																																			
<p>ASAC 55.4 DETAILSPUNAL LITERATURE CLASSIFICATION</p>																																																			

BC

B3
1

1ST AND 2ND ORDERS

PROCESSES AND PROPERTIES INDEX

Digestibility coefficients of some fodder substitutes for horses.
J. Skulmowski (*Ann. Univ. M. Curie-Skłodowska*, 1950, 8, DD, 13-27).—The digestibility coeff. of diets consisting of oats, hay, straw, and blood- or meat-meal cake are 7-15% lower than those calculated from Kellner's tables. The fodder rations should be supplemented accordingly.
R. Truscek.

ASB-5LA METALLURGICAL LITERATURE CLASSIFICATION

RECORD #1

RECORD #2

RECORD #3

RECORD #4

RECORD #5

RECORD #6

RECORD #7

RECORD #8

RECORD #9

RECORD #10

RECORD #11

RECORD #12

RECORD #13

RECORD #14

RECORD #15

RECORD #16

RECORD #17

RECORD #18

RECORD #19

RECORD #20

RECORD #21

RECORD #22

RECORD #23

RECORD #24

RECORD #25

RECORD #26

RECORD #27

RECORD #28

RECORD #29

RECORD #30

RECORD #31

RECORD #32

RECORD #33

RECORD #34

RECORD #35

RECORD #36

RECORD #37

RECORD #38

RECORD #39

RECORD #40

RECORD #41

RECORD #42

RECORD #43

RECORD #44

RECORD #45

RECORD #46

RECORD #47

RECORD #48

RECORD #49

RECORD #50

RECORD #51

RECORD #52

RECORD #53

RECORD #54

RECORD #55

RECORD #56

RECORD #57

RECORD #58

RECORD #59

RECORD #60

RECORD #61

RECORD #62

RECORD #63

RECORD #64

RECORD #65

RECORD #66

RECORD #67

RECORD #68

RECORD #69

RECORD #70

RECORD #71

RECORD #72

RECORD #73

RECORD #74

RECORD #75

RECORD #76

RECORD #77

RECORD #78

RECORD #79

RECORD #80

RECORD #81

RECORD #82

RECORD #83

RECORD #84

RECORD #85

RECORD #86

RECORD #87

RECORD #88

RECORD #89

RECORD #90

RECORD #91

RECORD #92

RECORD #93

RECORD #94

RECORD #95

RECORD #96

RECORD #97

RECORD #98

RECORD #99

RECORD #100

SKULMOWSKI, J.

Chemical Abstracts
May 25, 1954
Biological Chemistry

(2)

The digestibility coefficients of some feeds used in feeding pigs. R. Drzas and J. Skulmowski. *Roczniki Nauk Rolniczych* 66, Ser. B, *Zootech.* No. 2, 83-102 (in English, 103-4) (1953).—Digestibility coeffs. (1) for pigs were detd. for wheat, barley, potatoes, beets, fish and blood meal, etc., and compared with data found in the literature. The difference between the data was so great that it was impossible to adopt any definite av. values for nutritive values, or for rationing. I is given for total dry substance, total org. matter, proteins, fat, N-free ext., and fiber for each of the feeds evaluated. Richard Ehrlich

SKULMOWSKI, J.

Chemical Abstracts
May 25, 1954
Biological Chemistry

(2)
Digestibility coefficients of some poultry feeds. J. Skulmowski and W. Tytlo. *Roczniki Nauk Rolniczych* 66, Ser. B, Zootech. No. 3, 21-41 (in English, 41-2) (1953).—
Digestibility coeffs. (I) of certain poultry feeds were detd. I for total dry substance for the grain of wheat was 75.85, barley 70.36, corn 82.42, oats 60.20, rye 78.48, soybeans 22.34, meat and bone meal 36.49, blood meal 64.73, fish meal 35.41, and a mash composed of crushed grain of oats, wheat, corn, barley bran, and of meat and bone meal 66.66%. The I values for total proteins, org. matter, ether ext., N-free ext., and fibers are given for each of the feeds. I values can be applied in establishing the feed rations for poultry. The feeding value of the mash was very high. Richard Ehrlich

SKULMOWSKI, J.

P. U. L. 83

✓ Determination of amino acids in hydrolyzates of hays of *Medicago sativa*, *Ononis sativa*, *Trifolium pratense*, *Vicia faba minor*, and *Lupinus luteus* by the paper chromatography method. Tadeusz Mierzejewski and Józef Skulmowski. *Ann. Univ. Mariae Curie-Skłodowska*, Lublin, **Polonia**, Sect. DD, 8, 387-98(1953/54)(English summary).

—The hays of these 5 plants were hydrolyzed with HCl at 120° for 6 hrs., then given the same treatment for 24 hrs. at room temp. The hydrolyzates were kept at -4° and a two-dimensional paper chromatography was undertaken on Whatmann No. 1 and 41 paper of 26 × 28 cm. size; 0.016 ml. of hydrolyzate was used each time. In one direction the chromatogram was developed with $C_6H_5OH-H_2O$ 7:3, in the other one with $PrOH-H_2O$ 7:3. It was possible in this way to identify clearly, in addn. to one spot which could not be ascribed to any known substance, the following acids: leucine (I), isoleucine (II), methionine (III), valine (IV), phenylalanine (V), proline, tryptophan (VI), tyrosine, alanine, threonine, lysine, glycerol, serine, glutamic acid, aspartic acid, asparagine, cystine (VII), arginine (VIII), and histidine (IX). I and II and also II, IV, and V appeared together on the same spot in the two-dimensional chromatogram, but could be easily sep'd.: I from II with $BuOH-AcOH-H_2O$, II, IV, and V from each other with $BuOH-MeOH-H_2O$, III from V with $C_6H_5OH-H_2O-NH_3$. III, VI, VII, VIII, and IX are present in minute amts. only. From the size and colors which the spots show with ninhydrin *V. faba minor* and *O. sativa* form one group of plants; the other 3 belong to a sep. group. Werner Jacobson

Skulmowski, J.

Suitability of the indicator method for determining digestibility coefficients with pigs: digestibility of fish pastes and mushroom meal. J. Krupka, Z. Piaszczyński and J. Skulmowski (*Roczn. Nauk rol.*, 1955, 69, B, 195—210).—In digestibility trials with pigs Cr_2O_3 is preferable to SiO_2 as an indicator substance. For Norwegian fish pastes and mushroom meal (*Cantharellus cibarius*), respectively, the digestibility data were: org. matter, 83.4, 89.6; crude protein, 78.7, 53.9; true protein, 89.5, 66.3; fibre, —, 85.0; N-free extractives, —, 91.7%.
A. G. POLLARD.

SKULMOWSKI, J.

POLAND/Farm Animals. General Problems

Q-1

Abs Jour : Ref Zhur - Biol., No 8, 1958, No 35613

Author : ~~Skulmowski J.~~

Inst : Not Given

Title : Antibiotics in Animal Feeding

Orig Pub : Pustopy nauk roln., 1957, 3, No 5, 63-82

Abstract : The author presents a review on the use and stimulating action of antibiotics in swine breeding, poultry raising, and cattle breeding. A separate chapter is devoted to the effect of antibiotics upon the microflora of the stomach. A bibliography of 100 items is added.

Card : 1/1

SKULMOWSKI, Jozef

SURNAME, Given Names

Country: Poland

Academic Degrees:

Affiliation:

Source: Warsaw, Medycyna Weterynaryjna, Vol XVII, No 8, August 1961, pp 471-474

Data: "Nutritional Values of Mixed Feed Concentrates Produced by the Fodder Industry."

Authors:

SKULMOWSKI, Jozef; Prof. dr.; Director of the Biochemistry Department (Zaklad Biochemii), Veterinary Institute (Instytut Weterynarii), Pulawy; and p f the Department of Physiological Chemistry (Zaklad Chemii Fizjologicznej), Higher Agricultural School (WSR-- Wyzsza Szkola Rolnicza) Lublin.

DRZAS, Edward

PLISZCZYNSKI, Zbigniew

870 981643

POLAND

OLSKO-GRUNDROECK, Jadwiga, Dc. and RASZEWSKA, Helena of the Department of Biochemistry (Zaklad Biochemii) (Director: Prof. Dr. J. SKULMOWSKI) and of the Department of Poultry Diseases (Zaklad Chorob Drobiu) (Director: Docent, Dr. K. JARSKI), both of the Veterinary Institute (Instytut Weterynaryjny) in Poland.

"Variation of Serum Protein Fractions Level and HI and SN Titres According to Mode of Inoculation with 'F' Vaccine Against Newcastle Disease."

Warsaw-Lublin, Medycyna Weterynaryjna, Vol 13, No 9, Sep 23, pp. 11-137.

Abstract: [Authors' English summary modified]. Materials, methods, and results are given on the above study in 20 hens. Inoculation was made via the nose, intramuscular, and into the wing. Results are discussed indicating that immunity, as demonstrated by SN test, is probably due to alpha globulin fraction, and that vaccination via nose is most effective. Of the ten references, four are English.

PROCESS AND PROPERTIES INDEX	
<p>A SKULOV, N. S. N</p> <p>3C-272. Investigation of Constant of Anisotropy of Energy of Ternary Alloys of the System Ni-Cu-Mo. (In Russian.) N. S. Skulov, O. I. Blokhina, K. M. Bol'shova. Zhurnal Tekhnicheskoi Fiziki (Journal of Technical Physics), v. 19, Aug. 1949, p. 865-870.</p> <p>The influence of each element was investigated for multicomponent systems of Ni with nonferromagnetic elements. On the basis of previous investigations of binary alloys of Ni-Cu and Ni-Mo, the question of how the constant of anisotropy decreases in Ni-Cu-Mo alloys with increase of amount of Cu and Mo was solved. As a result, it is assumed that the above relationship has a universal character.</p>	
<p>ASB-5L-6 METALLURGICAL LITERATURE CLASSIFICATION</p>	

MALEK, P.; CERVINKA, F.; SKULOVA, M.; HOUSKA, O.

Epidemiologic studies on pemphigus mastitis strain of Staphylococcus; maternal infection and its relation to newborn infant and environment. Cesk. gyn. 18 no.4:299-306 Aug 1953. (CIML 25:4)

1. Of the Institute of Experimental Surgery (Director--Docent B. Spacek, M.D.) and of the Institute of Mother and Child (Director--Prof. J. Trapl, M.D.), Prague. 2. Epidemiological study.

MALEK, Prokop; Technicka spoluprace: JEHLICKOVA, M.; SKULOVA, M.;
STYS, V.

Dynamics of circulation of substances in the organism during
shock. Cas. lek. cesk. 95 no.17:449-454 27 Apr 56.

1. Ustav klinicke a experimentalni chirurgie, Praha.

(SHOCK, experimental

hemodynamics of procaine penicillin in animals. (Cz))

(PENICILLIN, derivatives

procaine penicillin, hemodynamics in expr. shock in
animals. (Cz))

VOL', L.M.; SKULOVICH, L.I.

Modernization of the crucible electric furnace for melting aluminum and its alloys with Nichrome-band heaters. Biul. tekhn.-ekon.inform. Gos. nauch.-issl. nauch. i tekhn. inform. 18 no.6:36-37 Je '65. (MIRA 18:7)

BIALAS, Zabuwa, A.; SMULSKA, E.; WALACH, Z.

Relative line strengths in the doublets of the spark spectra of Mg II, Ca II, Sr II, and Ba II. *Acta physica Pol* 26 no.2:175-183 '64.

1. Institute of Physics of the Jagiellonian University, Krakow.

VERZHBINSKAYA, N.A.; SKUL'SKAYA, G.A.

Riboflavin in the brain of vertebrates. Dokl. AN SSSR 110
no.6:1034-1037 0 '56. (MLRA 10:2)

1. Institut fiziologii imeni I.P. Pavlova Akademii nauk SSSR.
Predstavleno akademikom V.A. Engel'gardtom.
(Riboflavin) (Brain)

SKUL'SKAYA, G. A.

✓ Riboflavin in the brain of the vertebrates. N. A. Verzhbinskaya and G. A. Skul'skaya. *Doklady Akad. Nauk S.S.S.R.* 110, 1034-7 (1956). Typical analyses are given for riboflavin in the brain, heart, liver, kidney, and muscle tissues of 14 species of fish, frog, turtle, hen, grackle, bluebird, guinea pig, rabbit, hedgehog, and cat. In the vertebrate brain the content of riboflavin is relatively const. at about 2 g./l.; the exceptions are beluga fish (7-12) and frog (5-6.9); there is no definite progression in the evolutionary scale. Cytochrome activity in the brain increases steadily with evolutionary position from Q_{01} of 10 for lower fish to 55 in white rat; the ratio of cytochrome to riboflavin also shows a general progression in this order. Individual variations are discussed. G. M. Kosolapoff

2

L 13382-66 EWT(1)/FS(v)-3 SCTB DD

ACC NR: AP6002682 SOURCE CODE: UR/0385/65/001/006/0571/0576

AUTHOR: Barbashova, Z. I.; Skul'skaya, G. A.; Grigor'yeva, G. I.;
Vasil'yeva, V. V.

34
B

ORG: Study Group on Resistance of the Institute of Evolutionary
Physiology and Biochemistry im. I. M. Sechenov, AN SSSR, Leningrad
(Gruppa po izucheniuyu rezistentnosti Instituta evolyutsionnoy
fiziologii i biokhimii AN SSSR)

TITLE: Investigation of some properties of actomyosin proteins in
normal and hypoxic-conditioned rats 2

SOURCE: Zhurnal evolyutsionnoy biokhimii i fiziologii, v. 1, no. 6,
1965, 571-576

TOPIC TAGS: hypoxia, ~~muscle~~ protein, ~~animal~~ muscle physiology, rat,
space chamber, myology

ABSTRACT: The physical and chemical properties of animal-muscle
protein were studied to determine whether a relationship exists
between prolonged hypoxia conditioning and the ability of muscle to
resist the damaging effect of certain agents, such as protein-denaturing
substances. Tests were performed on contractile protein (actomyosin
group), which comprises 40% of the total muscle protein. White rats
weighing 250—300 g received hypoxia conditioning consisting of daily

Card 1/2 UDC: 591.175.05.044.01:547.96+612.744.015.33.014.41+616-001.12

L 13382-66

ACC NR: AP6002682

4-hr stays in an altitude chamber under gradually decreasing pressure. Actomyosin was then extracted from the femur muscles of decapitated experimental and control animals, and tests (described in detail in the original article) were performed. Experimental results showed that the following properties of actomyosin from control and conditioned animals were identical: a) the amount of extractable actomyosin, b) the specific and characteristic viscosity of actomyosin, c) the content of sulfhydryl groups, and d) the resistance to urea denaturing. Thus, the nonspecifically increased resistance of the skeletal musculature of hypoxic-conditioned rats observed in previous experiments is not related to the properties of actomyosin studied. However, in hypoxic-conditioned animals, the rate of recovery of actomyosin viscosity after the influence of ATP was significantly greater than the recovery rate in control animals. The mechanism of this phenomenon is not yet understood. Orig. art. has: 2 figures and 1 table. [JS]

SUB CODE: 06/ SUBM DATE: 25Sep64/ ORIG REF: 012/ OTH REF: 004
ATD PRESS: 4/54

Card 2/2 *gc*

PROTSENKO, L.D.; SKUL'SKAYA, N.Ya.

Aryl diethylene triamides of thiophosphoric acid. Zhur.ob.khim.
33 no.7:2284-2287 J1 '63. (MIRA 16:8)
(Phosphoroamidothioic acid)

L 17952-65 EWT(m)/EPF(c)/EWP(j) Pc-4/Pr-4 RPL RM
 ACCESSION NR: AP5002563 S/0079/64/034/007/2233/2234

AUTHOR: Protsenko, L. D.; Skul'skaya, N. Ya.

TITLE: O-aryl-N,N,N',N' - diethylenediamides of thiophosphoric acid B

SOURCE: Zhurnal obshchey khimii, v. 34, no. 7, 1964, 2233-2234

TOPIC TAGS: phosphoric acid, organic synthetic process, chloride, phenol

Abstract: A series of six O-aryl-N,N,N',N' -diethylenediamides of thiophosphoric acid and chlorides of O-arylthiophosphoric acids were synthesized (Ar = p-CH₃C₆H₄, o-CH₃C₆H₄, p-ClC₆H₄, p-IC₆H₄, p-BrC₆H₄, and p-NO₂C₆H₄). The O-aryl-N,N,N',N' -diethylenediamides of thiophosphoric acid were colorless crystalline compounds or viscous light yellow oils. The initial chlorides of the corresponding O-arylthiophosphoric acids were produced by the action of phosphorus thiochloride on phenols in pyridine. Orig. art. has 2 tables.

ASSOCIATION: none

SUBMITTED: 17Jun63

ENCL: 00

SUB CODE: OC, GC

NO REF SOV: 000

OTHER: 002

JPRS

Card 1/1

PROTSENKO, L.D.; SKUL'SKAYA, N.Ya.

Reaction of arylenephosphoramides and aroxyethylenethio-
phosphoramides with chlorine and hydrogen chloride. Zhur. ob.
khim. 35 no.4:715-717 Ap '65. (MIRA 18:5)

MAKAR'YEVA, A.G.; SKUL'SKAYA, B.V. .

Mechanization of minor operations for cold gluing of shaped parts.
Der.prom.5 no.8:21 Ag '56. (MLRA 9:10)

1.Tallinskaya fanerno-mebel'naya fabrika.
(Talinn--Furniture industry) (Gluing)

SKUL'SKAYA, R.V., inzh.

Automatic three-spindle machines for drilling and counterboring
holes in bars. Der.prom. 7 no. 7:22 J1 '58. (MIRA 11:8)

1. Tallinskaya fanerno-mebel'naya fabrika.
(Drilling and boring machinery)

SKUL'SKAYA, R.V., inzh.

Polishing by textile wheels without using shellac. Der.prom. 7 no.9:25-26
S '58. (MIRA-11:11)

1. Tallinskaya fanerno-mebel'naya fabrika.
(Wood finishing)

SKUL'SKAYA, R.V.

Making furniture with bent and glued parts from veneer sheets.
Der. prom. 13 no.1:17-20 Ja '64. (MIRA 17:4)

1. Tallinskaya fanerno-mebel'naya fabrika.

KOVALEVSKAYA, I.L.; EPSHTEYN-LITVAK, R.V.; DMITRIYEVA-RAVIKOVICH, Ye.M.;
 KURNOSOVA, N.A.; SHCHEGLOVA, Ye.S.; FERDINAND, Ya.M.;
 KHCMIK, S.R.; MAKHLINOVSKIY, L.P.; PETROVA, S.S.;
 GOLUBOVA, Ye.Ye.; GONCHAROVA, Z.I.; SARMANEYEV, A.P.;
 SIZINTSEVA, V.P.; Primali uchastiye: MEDYUKHA, G.A.;
 OSOKINA, L.A.; RACHKOVSKAYA, Yu.K.; OSOVTSEVA, O.I.;
 DEDUSENKO, A.I.; KOVALEVA, P.S.; KARASHEVICH, V.P.;
 CHEBOTAREVICH, N.D.; CHIGIR', T.R.; SKUL'SKAYA, S.D.;
 KECHETZHIYEV, B.A.; DEMINA, A.S.; ZUS'MAN, R.T.; YESAKOV, P.I.;
 SYSOYEVA, Z.A.; ZINOV'YEVA, I.S.; FAL'CHEVSKAYA, A.A.;
 DENISOVA, B.D.; TIMOFELEVA, R.G.; SYRKASOVA, A.V.;
 LYANTSMAN, S.G.

Reactivity and immunological and epidemiological effectiveness
 of alcoholic typhoid and paratyphoid fever vaccines in school
 children. Zhur. mikrobiol., epid. i immun. 33 no.7:72-77
 J1 '62. (MIRA 17:1)

1. Iz Moskovskogo, Rostovskogo, Omskogo institutov epidemio-
 logii i mikrobiologii, Stavropol'skogo instituta vaktsin i
 syvorotok i Ministerstva zdravookhraneniya RSFSR. 2. Rostovski
 institut epidemiologii i mikrobiologii (for Kovaleva).
3. Stavropol'skiy institut vaktsin i syvorotok (for Sysoyeva).
4. Kuybyshevskiy institut epidemiologii i mikrobiologii (for
 Zinov'yeva). 5. Saratovskaya gorodskaya sanitarno-epidemiolo-
 gicheskaya stantsiya (for Lyantsman).

FERDINAND, Ya.M. (Rostov-na-Donu); Prinimali uchastiye: MARISOVA, A.P.;
BRAYNINA, R.A.; MARGULIS, L.A.; MYASNENKO, A.M.; KOVALEVSKAYA,
I.L.; TELESHEVSKAYA, E.A.; SOBOLEVA, S.V.; KALININA, K.I.;
KOVALEVA, N.S.; IVANOVA, M.K.; ARENDER, B.A.; KUCHERENKO, R.A.;
MANATSKOVA, K.S.; OLEYNIKOVA, L.T.; KIBARDINA, Yu.A.;
GRIGOR'YEVA, K.S.; SEMENIKHINA, L.G.; CHERNYKH E.I.; DOROFYEVA,
V.M.; SHEVCHENKO, Ye.N.; ABRAMOVA, O.K.; SKUL'SKAYA, S.D.;
PETROVA, Z.I.; MAKHLINOVSKIY, L.I.; KUZ'MINA, A.I.; AL'TMAN, R.Sh.;
MARDERER, R.G.; YENGALYCHEVSKAYA, L.N.; CHIRKOVA, M.N.; TERESHCHENKO,
N.I.; SHELKOVNIKOVA, M.A.; PROKOPENKO, V.V.; BEKLEMESHEVA, Ye.A.;
BARANOVA, T.V.

Effectiveness of specific prophylaxis with alcohol divaccine
against typhoid and paratyphoid B fever in school-age children.
Zhur. mikrobiol., epid. i immun. 41 no.1:23-27 Ja '64.

(MIRA 18:2)

Country	:	Poland	G-3
Category	:		
Abs. Jour	:		45987
Author	:	Jerznanowska, J. and Skulski, J.	
Institut.	:	Not given	
Title	:	Dimerization and Polymerization Reactions of Euparin	
Orig Pub.	:	Roczniki Chem, 32, No 3, 471-433 (1951)	
Abstract	:	<p>The substance (I) formed by the dimerization of euparin (2-isopropenylcoumarone) (II) in dil alcoholic solutions of mineral acids (see preceding communication, RZhKhim, No 23, 1955, 55256), absorbs 3 mols H_2 on hydrogenation over Pt (from PtO_2) in glacial CH_3COOH and adds 3 mols perbenzoic acid in chloroform solution. On refluxing for 19 hrs in C_6H_6 with maleic anhydride, I forms the adduct $C_{30}H_{26}O_9$, mp 120°, later 250° [sic] (decomp; from benzene), and on</p>	

Card: 1/5

Country	:	Poland	G-3
Category	:		
Abstr. Jour	:		45987
Author	:		
Institut.	:		
Title	:		
Orig Pub.	:		
Abstract	:	<p>OH₂ group at the double bond in I. When a 10% benzene solution of II is left to stand in the presence of a 48% ether solution of BF₃ (several days at 20°), an amorphous polymer (III) with no sharply defined mp is formed: judging from the molecular weight this polymer appears to be a mixture of tetramer and small quantities of pentamer. The ozonolysis of III gives 3H₂O. I does not polymerize with a mixture of I and II. Attempts to brominate, sulfonate, and nitrate II were unsuccessful.</p> <p style="text-align: right;">D. Vichovski</p>	

Card: 3/3

SKULSKI, L.; URBANSKI, T.

On azo dyes deriving from 4-(*ω* -nitrovinyl)-aniline. Bul Ac Pol
chim 6 no.5:293-298 '58. (EEAI 9:7)

1. Department of Organic Technology, Warsaw Technical University.
Communicated by T.Urbanski.
(Azo dyes) (Nitrovinylaniline)

SKULSKI, Lech, mgr. inż.; URBANSKI, Tadeusz

On the absorption spectra of azo dyes. I. Spectra of azo dyes
deriving from 4-hydroxy- and 4-acetoxiazobenzene. Roczniki chemii 34
no.1:141-157 '60. (EEAI 10:9)

1. Department of Organic Technology II, Institute of Technology,
Warsaw.

(Absorption spectra)	(Azo compounds)	(Azobenzene)
(Acetoxy group)	(Phenylazophenol)	

SKULSKI, Lech; URBANSKI, Tadeusz

Reactions and the absorption spectra of compounds derived from
w-nitrostyrene. I. On the preparation of 4-(β -nitrovinyl)- aniline.
Rocz chemii 34 no.1:283-287 '60. (EEAI 10:9)

1. Department of Organic Technology II, Institute of Technology,
Warsaw.

(Absorption spectra) (Nitrostyrene)
(Nitrovinylaniline)

SKULSKI, Lech; URBANSKI, Tadeusz

Reactions and absorption spectra of compounds deriving from *u*-nitro-styrene. Pt.2. On synthesis of azo dyes with a β -nitrovinyl group. Roczniki chemii 34 no.2:431-441 '60. (EEAI 10:1)

1. Katedra Technologii Organicznej II Politechniki, Warszawa.
(Absorption spectra) (Azo dyes)
(Nitrovinyl group) (Nitrostyrene)